

## MYCOTAXON

<http://dx.doi.org/10.5248/120.119>

Volume 120, pp. 119–125

April–June 2012

***Dissoconium proteae* newly recorded from China**

RONG ZHANG<sup>1</sup>, YONGNA MAO<sup>1</sup>, LU HAO<sup>1</sup>,  
HONGCAI CHEN<sup>2</sup>, GUANGYU SUN<sup>1\*</sup> & MARK L. GLEASON<sup>3</sup>

<sup>1</sup> State Key Laboratory of Crop Stress Biology in Arid Areas and College of Plant Protection

& <sup>2</sup> College of Life Sciences, Northwest A&F University, Yangling, Shaanxi 712100, P. R. China

<sup>3</sup> Department of Plant Pathology and Microbiology, Iowa State University, Ames, Iowa 50011, U.S.A

\* CORRESPONDENCE TO: [sgy@nwsuaf.edu.cn](mailto:sgy@nwsuaf.edu.cn)

**ABSTRACT** — We document the first report of *Dissoconium proteae* from China. This fungus was found on the cuticle of apple fruit collected from orchards in Shaanxi Province. Its morphology is compared with other *Dissoconium* species and a phylogenetic analysis based on ITS sequence is presented.

**KEY WORDS** — sooty blotch and flyspeck, taxonomy, internal transcribed spacer, rDNA

**Introduction**

Sooty blotch and flyspeck (SBFS) is a worldwide disease complex more than 60 fungi that blemish the cuticle of pome fruits (such as apple) in humid temperate regions and cause cosmetic damage entailing significant economic losses (Batzler et al. 2005). The common name “flyspeck” refers to species in the SBFS complex that appear on fruit surfaces as clusters of black, shiny, round to oval, flattened sclerotium-like bodies with no mycelial mat (Batzler et al. 2005).

The hyphomycete genus *Dissoconium* de Hoog et al. has lightly pigmented conidiophores that produce septate and aseptate conidia in a sympodial manner. The conidia are hyaline, one- or two-celled, and detach forcibly in pairs (Hoog et al. 1983).

Currently the genus includes nine species: *Dissoconium aciculare* de Hoog et al., *D. australiense* Crous & Summerell, *D. commune* Crous & Mansilla, *D. dekkeri* de Hoog & Hijwegen, *D. eucalypti* Crous & Carnegie, *D. mali* G.Y. Sun et al., *D. musae* Arzanlou & Crous, *D. proteae*, and *D. subuliphorum* (Matsush.) R.F. Castañeda (Arzanlou et al. 2008; Crous et al. 1999, 2004, 2007, 2008; Hoog et al. 1983, 1991; Zhang et al. 2007). Batzler et al. (2005) reported five putative *Dissoconium* species within the sooty blotch complex on apple fruit.

Recently, during investigations of SBFS of apple in China, we identified four fungal isolates that represent the first record of *Dissoconium proteae* from China based on ITS sequence analysis and morphological comparison.

**Materials & methods**

**ISOLATION.** In September 2008, apples with flyspeck signs were collected from orchards in Shaanxi Province. The thalli were transferred to a potato dextrose agar slant (potato 200 g, dextrose 20 g, agar 10 g; PDA) directly from the apple surface, and then cultured at 23±1°C in darkness (Sun et al. 2003). Colony description of the species was based on a 1-month-old pure culture. Morphological characteristics are described from specimens obtained from a cover slip inserted into the surface of the media. Voucher specimens and cultures are conserved in the Fungal Herbarium of Northwest A&F University, Yangling, China (HMUABO).

**DNA SEQUENCING.** Template DNA was extracted from fungal mycelium according to Li et al. (2011); primer pairs used for ITS amplification and sequencing were ITS1-F

TABLE 1. Sequences used in the phylogenetic analysis

| SPECIES                                                      | GENBANK CODE | REFERENCE                |
|--------------------------------------------------------------|--------------|--------------------------|
| <i>Dissoconium proteae</i> (mynbxd10)                        | HM070373     | This paper               |
| <i>D. proteae</i> (mynbxd2)                                  | HM070374     | This paper               |
| <i>D. proteae</i> (mynbxd1.1)                                | HM070375     | This paper               |
| <i>D. proteae</i> (mynxc1.1)                                 | HM070376     | This paper               |
| <i>D. aciculare</i>                                          | AY725519     | Crous et al. (2004)      |
|                                                              | AY725521     | Crous et al. (2004)      |
| <i>D. australiense</i>                                       | EF394854     | Crous et al. (2007)      |
| <i>D. eucalypti</i>                                          | EF394855     | Crous et al. (2007)      |
| <i>D. mali</i>                                               | EF627451     | Zhang et al. (2007)      |
| <i>D. musae</i>                                              | EU514226     | Arzanlou et al. (2008)   |
|                                                              | EU514225     | Arzanlou et al. (2008)   |
| <i>D. proteae</i>                                            | EU707897     | Crous et al. (2008)      |
| <i>Dissoconium</i> sp.                                       | AY598874     | Batzer et al. (2005)     |
|                                                              | AY598875     | Batzer et al. (2005)     |
|                                                              | AY598877     | Batzer et al. (2005)     |
|                                                              | FJ425205     | Díaz Arias et al. (2010) |
| <i>Mycosphaerella communis</i><br>(anam. <i>D. commune</i> ) | AY725544     | Crous et al. (2004)      |
|                                                              | DQ302950     | Crous et al. (2006)      |
| <i>M. lateralis</i> (anam.: <i>D. dekkeri</i> )              | EU301078     | Burgess et al. (2007)    |
| <i>Ramichloridium apiculatum</i>                             | EU041794     | Arzanlou et al. (2007)   |
|                                                              | EU041791     | Arzanlou et al. (2007)   |
| <i>R. cerophilum</i>                                         | EU041798     | Arzanlou et al. (2007)   |
| <i>R. indicum</i>                                            | EU041799     | Arzanlou et al. (2007)   |
| <i>Ramichloridium</i> sp.                                    | FJ425199     | Díaz Arias et al. (2010) |
| <i>Stenella araguata</i>                                     | AF362066     | Crous et al. (2001)      |

(Gardes & Bruns 1993) and ITS4 (White et al. 1990). Amplification was completed with the following cycling parameters: initial denaturation at 94°C for 3 min followed by 35 cycles of denaturation at 94°C for 30 s, annealing at 52°C for 30 s, and extension at 72°C for 10 min. The PCR products were sequenced by Organism Technology Co., Shanghai, China.

The ITS nucleotide sequences generated in this study were added to sequences downloaded from GenBank (TABLE 1) that had high similarity according to a BLAST search (National Center for Biotechnology Information's nucleotide blast program). Preliminary alignments were performed using CLUSTAL-X (1.83) (Thompson et al. 1997), imported into BioEdit version 5.0.9.1 (Hall 1999), and manually adjusted. Phylogenetic analysis of aligned DNA sequences was performed with PAUP version 4.0b10 for 32-bit Microsoft Windows (Swofford 2001). Heuristic searches were performed with 1000 random sequence additions. Clade stability was evaluated by 1000 bootstrap replications. Other measures for parsimony, including tree length, consistency index, retention index and rescaled consistency index (CI, RI and RC, respectively), were also calculated. *Stenella araguata* was used as the outgroup taxon.

### Results: DNA phylogeny

A multiple alignment of the rDNA-ITS was generated with 21 sequences obtained from GenBank plus the sequences of isolates mynxbd10, mynxbd2, mynxbd1.1 and mynxc1.1. An MP tree with 354 length (CI = 0.7232, RI = 0.8622, RC = 0.6235) was constructed (Fig. 1). Two major clades were resolved in the MP trees. One of the major clade with 100% bootstrap value contained three species in *Dissoconium*, *Mycosphaerella*, and *Ramichloridium*. The other major clade also had a bootstrap value of 100%. Our isolates and an authentic *Dissoconium proteae* isolate identified by Crous et al. (2008) fell within a single clade with 71% bootstrap support.

### Taxonomy

*Dissoconium proteae* Crous, Persoonia 20: 68. 2008.

FIG. 2

Mycelium external consisting of branched, septate, smooth, hyaline hyphae, 0.8–2.8 µm wide. Conidiophores 9–20 × 2.2–4.7 µm, solitary, arising from hyphae, subcylindrical, subulate, tapering to a bluntly rounded or truncate apex, straight to gently curved, smooth, hyaline, becoming medium pale brown with age, aseptate; conidiogenous loci terminal and lateral, visible as slightly thickened, darkened scars, 0.5 µm wide. Conidia 6.3–11.5 × 2.2–4 µm, solitary, hyaline to pale olivaceous, smooth, ellipsoid, non-constricted to slightly constricted at median septum, apex obtuse, base obconic-truncate, tapering pronounced at somewhat protruding hilum, unthickened, not darkened, 1 µm wide. Secondary conidia developing adjacent to primary conidia, hyaline to subhyaline, aseptate, ellipsoid, tapering prominently towards a protruding, truncate base, 4.3–8 × 1.6–3.8 µm; anastomosing with primary conidia after




FIG. 1 The majority consensus tree (length = 354, CI = 0.7232, RI = 0.8622, RC = 0.6235) derived from a heuristic search option in PAUP version 4.0b10 for 32-bit Microsoft Windows with 1000 randomizations of sequence input orders and 1000 bootstrap replications using the data set of ITS1, 5.8S and ITS2. Bootstrap values higher than 50% are indicated above or below the tree branches.

active discharge. In some cases the secondary conidia were observed to germinate.

**CULTURAL CHARACTERISTICS:** Colonies on PDA spreading, with sparse aerial mycelium and regular smooth margins; surface white or sienna, with patches of white and cinnamon; forming clusters of black sclerotia (remaining infertile) on PDA; reaching 15 mm diam after 1 mo.

**CHARACTERISTICS ON THE HOST:** On apple peel, the fungus produced dark, shiny, round to oval, slightly protuberant sclerotium-like bodies (FIG. 2A).

**SPECIMENS EXAMINED:** China: Shaanxi Province, Xianyang, Bin County, 35°04'N 108°09'E, alt. 1108 m, on *Malus ×domestica* (Rosaceae), 18 Sep. 2008, Y.N. Mao, HMUABO (the Fungal Herbarium of Northwest A&F University) 8885, 8877, 8866 (with dried culture), culture mynxbd10, mynxbd2, mynxbd1.1. China: Shaanxi Province, Xianyang, Xunyi County, 35°13'N 108°33'E, alt. 1300 m, On *Malus ×domestica*, 18 Sep. 2008, Y.N. Mao, HMUABO 8682 (with dried culture), culture mynxc1.1.




FIG. 2 *Dissoconium proteae* isolate mynboxd10. A. Signs on apple peel. B. Colony on PDA agar after 30 days. C-D. Conidia, conidiogenous cells, and hyphae. E. Conidiogenous cell. F. Conidium and secondary conidium. Bars: B=10 mm, C-F =10  $\mu$ m.

## Discussion

Of the more than 60 fungal species that blemish the surface of apples worldwide (Díaz Arias et al. 2010), many have been described quite recently. For example, Li et al. (2010) and Ma et al. (2010) reported three species of *Zygophiala* (*Z. cryptogama* and *Z. cylindrica*; *Z. qianensis*) associated with flyspeck on apples. Our report provides the first evidence that *Dissoconium proteae* can cause flyspeck on apple fruit. Crous et al. (2008) described *D. proteae* from leaves of *Protea* sp. (*Proteaceae*), causing a leaf spot.

Currently there are nine species in *Dissoconium*. Based on phylogenetic analysis of the ITS region and morphological characters of the anamorph, we identified four isolates as *Dissoconium proteae*. This is the first report of the species from apple, and a new record for China.

## Acknowledgments

This work was supported by National Natural Science Foundation of China (31170015, 31171797), the 111 Project from Education Ministry of China (B07049), Top Talent Project of Northwest A&F University and the earmarked fund for Modern Agro-industry Technology Research System (nycytx-08-04-04). The authors wish to thank Dr Eric H.C. McKenzie (Landcare Research, Auckland, New Zealand) and Professor Zhongyi Zhang (College of Plant Protection, Yunnan Agricultural University, Kunming, Yunnan, China) for reviewing the manuscript.

## Literature cited

- Arzanlou M, Groenewald JZ, Gams W, Braun U, Shin H-D, Crous PW. 2007. Phylogenetic and morphotaxonomic revision of *Ramichloridium* and allied genera. *Studies in Mycology* 58: 57–93. <http://dx.doi.org/10.3114/sim.2007.58.03>
- Arzanlou M, Groenewald JZ, Fullerton RA, Abeln ECA, Carlier J, Zapater M-F, Buddenhagen IW, Viljoen A, Crous PW. 2008. Multiple gene genealogies and phenotypic characters differentiate several novel species of *Mycosphaerella* and related anamorphs on banana. *Persoonia* 20: 19–37. <http://dx.doi.org/10.3767/003158508X302212>
- Batzer JC, Gleason ML, Harrington TC, Tiffany LH. 2005. Expansion of the sooty blotch and flyspeck complex on apples based on analysis of ribosomal DNA gene sequences and morphology. *Mycologia* 97: 1268–1286. <http://dx.doi.org/10.3852/mycologia.97.6.1268>
- Burgess TI, Pegg GS, Barber PA, Carnegie AJ, Hardy GESJ. 2007. <http://www.ncbi.nlm.nih.gov/nucleotide/162568812?report=genbank>
- Crous PW, Hong L, Wingfield MJ, Wingfield BD, Kang JC. 1999. *Uwebraunia* and *Dissoconium*, two morphologically similar anamorph genera with different teleomorph affinity. *Sydowia* 51: 155–166.
- Crous PW, Kang JC, Braun U. 2001. A phylogenetic redefinition of anamorph genera in *Mycosphaerella* based on ITS rDNA sequence and morphology. *Mycologia* 93: 1081–1101.
- Crous PW, Groenewald JZ, Mansilla JP, Hunter GC, Wingfield MJ. 2004. Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus*. *Studies in Mycology* 50: 195–214.
- Crous PW, Wingfield MJ, Mansilla JP, Alfenas AC, Groenewald JZ. 2006. Phylogenetic reassessment of *Mycosphaerella* spp. and their anamorphs occurring on *Eucalyptus*. II. *Studies in Mycology* 55: 99–131. <http://dx.doi.org/10.3114/sim.55.1.99>

- Crous PW, Summerell BA, Carnegie AJ, Mohammed C, Wingfield MJ, Himaman W, Groenewald JZ. 2007. Follicolous *Mycosphaerella* spp. and their anamorphs on *Corymbia* and *Eucalyptus*. *Fungal Diversity* 26:143–185.
- Crous PW, Summerell BA, Mostert L, Groenewald JZ. 2008. Host specificity and speciation of *Mycosphaerella* and *Teratosphaeria* species associated with leaf spots of *Proteaceae*. *Persoonia* 20:59–86. <http://dx.doi.org/10.3767/003158508X323949>
- Díaz Arias MM, Batzer JC, Harrington TC, Wong AW, Bost SC, Cooley DR, Ellis MA, Hartman JR, Rosenberger DA, Sundin GW, Sutton TB, Travis JW, Wheeler MJ, Yoder KS, Gleason ML. 2010. Diversity and biogeography of sooty blotch and flyspeck fungi on apple in the eastern and midwestern United States. *Phytopathology* 100: 345–355. <http://dx.doi.org/10.1094/PHYTO-100-4-0345>
- Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity for basidiomycetes– application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2:113–118. <http://dx.doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl Acids Symp Ser.* 41: 95–98.
- Hoog GS de, Oorschot CAN van, Hijwegen T. 1983. Taxonomy of the *Dactylaria* complex. II. *Dissoconium* gen. nov. and *Cordana* Preuss. *Proceedings van de Koninklijke Nederlandse Akademie van Wetenschappen, Section C.* 86: 197–206.
- Hoog GS de, Hijwegen T, Batenburg-van der Vegte WH. 1991. A new species of *Dissoconium*. *Mycological Research* 95: 679–682. [http://dx.doi.org/10.1016/S0953-7562\(09\)80814-9](http://dx.doi.org/10.1016/S0953-7562(09)80814-9)
- Li HY, Zhang R, Sun GY, Batzer JC, Gleason ML. 2010. New species and record of *Zygophiala* on apple fruit from China. *Mycological Progress* 9: 245–251. <http://dx.doi.org/10.1007/s11557-009-0633-1>
- Li HY, Sun GY, Batzer JC, Crous PW, Groenewald JZ, Karakaya A, Gleason ML. 2011. *Scleroramularia* gen. nov. associated with sooty blotch and flyspeck of apple and pawpaw from the Northern Hemisphere. *Fungal Diversity* 46:53–66. <http://dx.doi.org/10.1007/s13225-010-0074-9>
- Ma YQ, Zhang R, Sun GY, Zhu HX, Tang M, Batzer JC, Gleason ML. 2010. A new species of *Zygophiala* associated with the flyspeck complex on apple from China. *Mycological Progress* 9: 151–155. <http://dx.doi.org/10.1007/s11557-009-0635-z>
- Sun GY, Zhang R, Zhang Z, Zhang M. 2003. Isolation of sooty blotch and flyspeck fungi from apple surface by picking up the thalli. *Acta Phytopathology Sinica* 33: 479–480.
- Swofford DL. 2001. PAUP\*. Phylogenetic analysis using parsimony (\*and other methods) version 4.0 beta version. Sinauer Associates, Sunderland, Massachusetts, U.S.A.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The CLUSTAL-X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25: 4876–4882.
- White TJ, Bruns TD, Lee SB, Taylor JW. 1990. Analysis of phylogenetic relationships by amplification and direct sequencing of ribosomal RNA genes. 315–322, in: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds) PCR protocols: a guide to methods and applications. Academic, New York.
- Zhang R, Zhang Z, Zhang M, Sun GY, Gleason ML. 2007. A new species of *Dissoconium* from China colonizing apples. *Mycotaxon* 101: 165–172.